



Alcohol Facts and Statistics

Alcohol Use in the United States:

- » **Prevalence of Drinking:** According to the 2018 National Survey on Drug Use and Health (NSDUH), 86.3 percent of people ages 18 or older reported that they drank alcohol at some point in their lifetime; 70.0 percent reported that they drank in the past year; 55.3 percent reported that they drank in the past month.¹
- » **Prevalence of Binge Drinking and Heavy Alcohol Use:** In 2018, 26.5 percent of people ages 18 or older reported that they engaged in binge drinking in the past month; 6.6 percent reported that they engaged in heavy alcohol use in the past month.¹ (See sidebar on page 2 for definitions of binge drinking and heavy alcohol use.)



Alcohol Use Disorder (AUD) in the United States:

- » **Adults (ages 18+):** According to the 2018 NSDUH, 14.4 million adults ages 18 and older² (5.8 percent of this age group³) had AUD. This includes 9.2 million men² (7.6 percent of men in this age group³) and 5.3 million women² (4.1 percent of women in this age group³).
 - About 7.9 percent of adults who had AUD in the past year received treatment. This includes 8.0 percent of males and 7.7 percent of females with AUD in this age group.⁴
- » **Youth ages 12 to 17:** According to the 2018 NSDUH, an estimated 401,000 adolescents ages 12–17² (1.6 percent of this age group³) had AUD. This number includes 173,000 males² (1.4 percent of males in this age group³) and 227,000 females² (1.9 percent of females in this age group³).
 - About 5.0 percent of youth who had AUD in the past year received treatment. This includes 5.6 percent of males and 4.6 percent of females with AUD in this age group.⁴

Alcohol-Related Deaths:

- » An estimated 88,000⁵ people (approximately 62,000 men and 26,000 women⁵) die from alcohol-related causes annually, making alcohol the third leading preventable cause of death in the United States. The first is tobacco, and the second is poor diet and physical inactivity.⁶
- » In 2014, alcohol-impaired driving fatalities accounted for 9,967 deaths (31 percent of overall driving fatalities).⁷



Economic Burden:

- » In 2010, alcohol misuse cost the United States \$249.0 billion.⁸
- » Three-quarters of the total cost of alcohol misuse is related to binge drinking.⁸

Global Burden:

- » In 2012, 3.3 million deaths, or 5.9 percent of all global deaths (7.6 percent for men and 4.0 percent for women), were attributable to alcohol consumption.⁹
- » In 2014, the World Health Organization reported that alcohol contributed to more than 200 diseases and injury-related health conditions, most notably DSM–IV alcohol dependence (see sidebar), liver cirrhosis, cancers, and injuries.¹⁰ In 2012, 5.1 percent of the burden of disease and injury worldwide (139 million disability-adjusted life-years) was attributable to alcohol consumption.⁹
- » Globally, alcohol misuse was the fifth leading risk factor for premature death and disability in 2010. Among people between the ages of 15 and 49, it is the first.¹¹ In the age group 20–39 years, approximately 25 percent of the total deaths are alcohol attributable.¹²

Family Consequences:

- » More than 10 percent of U.S. children live with a parent with alcohol problems, according to a 2012 study.¹³

Underage Drinking:

- » **Prevalence of Underage Alcohol Use:**
 - **Prevalence of Drinking:** According to the 2018 NSDUH, 29.8 percent of 15-year-olds report that they have had at least 1 drink in their lives.¹⁴ About 7.1 million people ages 12–20¹⁵ (18.8 percent of this age group¹⁶) reported drinking alcohol in the past month (18.2 percent of males and 19.5 percent of females¹⁶).

Definitions

Alcohol Use Disorder (AUD): AUD is a chronic relapsing brain disease characterized by an impaired ability to stop or control alcohol use despite adverse social, occupational, or health consequences. AUD can range from mild to severe, and recovery is possible regardless of severity. The fourth edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM–IV)*, published by the American Psychiatric Association, described two distinct disorders—alcohol abuse and alcohol dependence—with specific criteria for each. The fifth edition, *DSM–5*, integrates the two DSM–IV disorders, alcohol abuse and alcohol dependence, into a single disorder called alcohol use disorder, or AUD, with mild, moderate, and severe subclassifications.

Binge Drinking:

- » NIAAA defines binge drinking as a pattern of drinking that brings blood alcohol concentration (BAC) levels to 0.08 g/dL. This typically occurs after 4 drinks for women and 5 drinks for men—in about 2 hours.³⁰
- » The Substance Abuse and Mental Health Services Administration (SAMHSA), which conducts the annual National Survey on Drug Use and Health (NSDUH), defines binge drinking as 5 or more alcoholic drinks for males or 4 or more alcoholic drinks for females on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past month.³¹

Heavy Alcohol Use: SAMHSA defines heavy alcohol use as binge drinking on 5 or more days in the past month.

Patterns of Drinking Associated with AUD: Binge drinking and heavy alcohol use can increase an individual's risk of AUD. The U.S. Dietary Guidelines recommend that if alcohol is consumed, it should be in moderation, defined as up to one drink per day for women and up to two drinks per day for men, and only by adults of legal drinking age. Some individuals should avoid alcohol completely.

Alcohol-Impaired-Driving Fatality: A fatality in a crash involving a driver or motorcycle rider (operator) with a BAC of 0.08 g/dL or greater.

Disability-Adjusted Life-Years (DALYs): A measure of years of life lost or lived in less than full health.

Underage Drinking: Alcohol use by anyone under the age of 21. In the United States, the legal drinking age is 21.



NIH . . . Turning Discovery Into Health®

National Institute on Alcohol Abuse and Alcoholism

<https://www.niaaa.nih.gov> • 301-443-3860

- **Prevalence of Binge Drinking:** According to the 2018 NSDUH, approximately 4.3 million people¹⁵ (about 11.4 percent¹⁶) ages 12–20 (11.3 percent of males and 11.4 percent of females¹⁶) reported binge drinking in the past month.
- **Prevalence of Heavy Alcohol Use:** According to the 2018 NSDUH, approximately 861,000 people¹⁵ (about 2.3 percent¹⁶) ages 12–20 (2.6 percent of males and 1.9 percent of females¹⁶) reported heavy alcohol use in the past month.

» **Consequences of Underage Alcohol Use:**

- Research indicates that alcohol use during the teenage years could interfere with normal adolescent brain development and increase the risk of developing AUD. In addition, underage drinking contributes to a range of acute consequences, including injuries, sexual assaults, and even deaths—including those from car crashes.¹⁷

Alcohol and College Students:

» **Prevalence of Alcohol Use:**

- **Prevalence of Drinking:** According to the 2018 NSDUH, 54.9 percent of full-time college students ages 18–22 drank alcohol in the past month compared with 44.6 percent of other persons of the same age.¹⁸
- **Prevalence of Binge Drinking:** According to the 2018 NSDUH, 36.9 percent of college students ages 18–22 reported binge drinking in the past month compared with 27.9 percent of other persons of the same age.¹⁸
- **Prevalence of Heavy Alcohol Use:** According to the 2018 NSDUH, 9.6 percent of college students ages 18–22 reported heavy alcohol use in the past month compared with 6.9 percent of other persons of the same age.¹⁸

» **Consequences—Researchers estimate that each year:**

- 1,825 college students between the ages of 18 and 24 die from alcohol-related unintentional injuries, including motor-vehicle crashes.¹⁹
- 696,000 students between the ages of 18 and 24 are assaulted by another student who has been drinking.²⁰
- 97,000 students between the ages of 18 and 24 report experiencing alcohol-related sexual assault or date rape.²⁰
- Roughly 20 percent of college students meet the criteria for AUD.²¹
- About 1 in 4 college students report academic consequences from drinking, including missing class, falling behind in class, doing poorly on exams or papers, and receiving lower grades overall.²²

Alcohol and Pregnancy:

- » The prevalence of Fetal Alcohol Syndrome (FAS) in the United States was estimated by the Institute of Medicine in 1996 to be between 0.5 and 3.0 cases per 1,000.²³
- » More recent reports from specific U.S. sites report the prevalence of FAS to be 2 to 7 cases per 1,000, and the prevalence of Fetal Alcohol Spectrum Disorders (FASD) to be as high as 20 to 50 cases per 1,000.^{24,25}



Alcohol and the Human Body:

- » In 2015, of the 78,529 liver disease deaths among individuals ages 12 and older, 47.0 percent involved alcohol. Among males, 49,695 liver disease deaths occurred and 49.5 percent involved alcohol. Among females, 28,834 liver disease deaths occurred and 43.5 percent involved alcohol.²⁶
- » Among all cirrhosis deaths in 2013, 47.9 percent were alcohol related. The proportion of alcohol-related cirrhosis was highest (76.5 percent) among deaths of persons ages 25–34, followed by deaths of persons ages 35–44, at 70.0 percent.²⁷
- » In 2009, alcohol-related liver disease was the primary cause of almost 1 in 3 liver transplants in the United States.²⁸
- » Drinking alcohol increases the risk of cancers of the mouth, esophagus, pharynx, larynx, liver, and breast.²⁹

For more information, please visit: <https://www.niaaa.nih.gov>

- 1 Substance Abuse and Mental Health Services Administration (SAMHSA). 2018 National Survey on Drug Use and Health (NSDUH). Table 2.1B—Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group: Percentages, 2017 and 2018. Available at: <https://www.samhsa.gov/data/sites/default/files/cbhsq-reports/NSDUHDetailedTabs2018R2/NSDUHDetTabsSect2pe2018.htm#tab2-1b>. Accessed 12/2/19.
- 2 SAMHSA. 2018 National Survey on Drug Use and Health (NSDUH). Table 5.4A—Alcohol Use Disorder in Past Year among Persons Aged 12 or Older, by Age Group and Demographic Characteristics: Numbers in Thousands, 2017 and 2018. Available at: <https://www.samhsa.gov/data/sites/default/files/cbhsq-reports/NSDUHDetailedTabs2018R2/NSDUHDetTabsSect5pe2018.htm#tab5-4a>. Accessed 12/2/19.
- 3 SAMHSA. 2018 National Survey on Drug Use and Health (NSDUH). Table 5.4B—Alcohol Use Disorder in Past Year among Persons Aged 12 or Older, by Age Group and Demographic Characteristics: Percentages, 2017 and 2018. Available at: <https://www.samhsa.gov/data/sites/default/files/cbhsq-reports/NSDUHDetailedTabs2018R2/NSDUHDetTabsSect5pe2018.htm#tab5-4b>. Accessed 12/2/19.
- 4 SAMHSA. Population prevalence estimates (%) are weighted by the person-level analysis weight and derived from the 2018 NSDUH public-use data file, defining “any treatment” as treatment or counseling designed to help reduce or stop alcohol use, including detoxification and any other treatment for medical problems associated with alcohol use, as well as defining AUD as alcohol abuse or alcohol dependence according to the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders*. National Survey on Drug Use and Health (NSDUH) 2018 (NSDUH-2019) Public-Use File Dataset. Available at: <https://www.datafiles.samhsa.gov/study/national-survey-drug-use-and-health-nsduh-2018-nid18757>. Accessed 11/15/19.
- 5 Centers for Disease Control and Prevention (CDC). *Alcohol and Public Health: Alcohol-Related Disease Impact (ARDI). Average for United States 2006–2010 Alcohol-Attributable Deaths Due to Excessive Alcohol Use*. Available at: https://nccd.cdc.gov/DPH_ARDI/Default/Report.aspx?T=AAM&P=f6d7eda7-036e-4553-9968-9b17ffad620e&R=d7a9b303-48e9-4440-bf47-070a4827e1fd&M=8E1C5233-5640-4EE8-9247-1ECA7DA325B9&F=&D=-. Accessed 1/18/17.
- 6 Mokdad, A.H.; Marks, J.S.; Stroup, D.F.; and Gerberding, J.L. Actual causes of death in the United States 2000. [Published erratum in: JAMA 293(3):293–294, 298] *JAMA: Journal of the American Medical Association* 291(10):1238–1245, 2004. PMID: 15010446
- 7 National Center for Statistics and Analysis. *2014 Crash Data Key Findings (Traffic Safety Facts Crash Stats*. Report No. DOT HS 812 219). Washington, DC: National Highway Traffic Safety Administration, 2015. Available at: <https://crashstats.nhtsa.dot.gov/Api/Public/ViewPublication/812219>. Accessed 9/19/16.
- 8 Sacks, J.J.; Gonzales, K.R.; Bouchery, E.E.; et al. 2010 national and state costs of excessive alcohol consumption. *American Journal of Preventive Medicine* 49(5):e73–e79, 2015. PMID: 26477807
- 9 World Health Organization (WHO). *Global Status Report on Alcohol and Health*. p. XIV. 2014 ed. Available at: http://www.who.int/substance_abuse/publications/global_alcohol_report/msb_gsr_2014_1.pdf?ua=1. Accessed 1/18/17.
- 10 World Health Organization (WHO). *Global Status Report on Alcohol and Health*. p. XIII. 2014 ed. Available at: http://www.who.int/substance_abuse/publications/global_alcohol_report/msb_gsr_2014_1.pdf?ua=1. Accessed 1/18/17.
- 11 World Health Organization (WHO). *Global Status Report on Alcohol and Health*. p. 57. 2014 ed. Available at: http://www.who.int/substance_abuse/publications/global_alcohol_report/msb_gsr_2014_1.pdf?ua=1. Accessed 6/15/17.
- 12 World Health Organization (WHO). *Alcohol*. 2015. Available at: <http://www.who.int/mediacentre/factsheets/fs349/en/#>. Accessed 9/19/16.
- 13 Substance Abuse and Mental Health Services Administration (SAMHSA). *Data Spotlight: More than 7 Million Children Live with a Parent with Alcohol Problems*, 2012. Available at: <https://www.samhsa.gov/data/sites/default/files/Spot061ChildrenOfAlcoholics2012/Spot061ChildrenOfAlcoholics2012.pdf>. Accessed 12/2/19.
- 14 SAMHSA. Population prevalence estimates (%) are weighted by the person-level analysis weight and derived from the 2018 NSDUH public-use data file. Lifetime alcohol use is defined as ever, even once, having had a drink of any type of alcoholic beverage (a can or bottle of beer, a glass of wine or a wine cooler, a shot of distilled spirits, or a mixed drink with distilled spirits in it), not counting a sip or two from a drink. National Survey on Drug Use and Health (NSDUH) 2018 (NSDUH-2018) Public-Use File Dataset. Available at: <https://www.datafiles.samhsa.gov/study-dataset/national-survey-drug-use-and-health-2018-nsduh-2018-ds0001-nid18758>. Accessed 11/18/19.
- 15 SAMHSA. 2018 National Survey on Drug Use and Health. Table 7.16A—Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Gender: Numbers in Thousands, 2002-2018. Available at: <https://www.samhsa.gov/data/sites/default/files/cbhsq-reports/NSDUHDetailedTabs2018R2/NSDUHDetTabsSect7pe2018.htm#tab7-16a>. Accessed 12/2/19.
- 16 SAMHSA. 2018 National Survey on Drug Use and Health. Table 7.16B—Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Detailed Age Category: Percentages, 2017 and 2018. Available at <https://www.samhsa.gov/data/sites/default/files/cbhsq-reports/NSDUHDetailedTabs2018R2/NSDUHDetTabsSect7pe2018.htm#tab7-16b>. Accessed 12/2/19.
- 17 National Institute on Alcohol Abuse and Alcoholism (NIAAA). *Alcohol Alert*, No. 67, “Underage Drinking,” 2006. Available at: <https://pubs.niaaa.nih.gov/publications/AA67/AA67.htm>. Accessed 9/19/16.



NIH . . . Turning Discovery Into Health®

National Institute on Alcohol Abuse and Alcoholism

<https://www.niaaa.nih.gov> • 301-443-3860

- ¹⁸ SAMHSA. The estimates are weighted by the person-level analysis weight and derived from the 2018 NSDUH public-use data file. Past-month alcohol use: a drink of an alcoholic beverage (a can or bottle of beer, a glass of wine or a wine cooler, a shot of distilled spirits, or a mixed drink with distilled spirits in it), not counting a sip or two from a drink in the past 30 days. Past-month binge alcohol use: 5 or more drinks on the same occasion for males or 4 or more drinks on the same occasion for females on at least 1 day in the past 30 days. Past-month heavy alcohol use: 5 or more drinks on the same occasion for males or 4 or more drinks on the same occasion for females on each of 5 or more days in the past 30 days. Full-time college students: full-time students ages 18 to 22 enrolled in school and at college level. Other persons aged 18–22: those not enrolled in school, enrolled in college part-time, enrolled in other grades either full- or part-time, or enrolled with no other information available. National Survey on Drug Use and Health (NSDUH) 2018 (NSDUH-2018) Public-Use File Dataset. Available at: <https://www.datafiles.samhsa.gov/study/national-survey-drug-use-and-health-nsduh-2018-nid18757>. Accessed 11/15/19.
- ¹⁹ Hingson, R.W.; Zha, W.; and Weitzman, E.R. Magnitude of and trends in alcohol-related mortality and morbidity among U.S. college students ages 18–24, 1998–2005. *Journal of Studies on Alcohol and Drugs* (Suppl. 16):12–20, 2009. PMID: 19538908.
- ²⁰ Hingson, R.; Heeren, T.; Winter, M.; et al. Magnitude of alcohol-related mortality and morbidity among U.S. college students ages 18–24: Changes from 1998 to 2001. *Annual Review of Public Health* 26:259–279, 2005. PMID: 15760289
- ²¹ Blanco, C.; Okuda, M.; Wright, C. et al. Mental health of college students and their non-college-attending peers: Results from the National Epidemiologic Study on Alcohol and Related Conditions. *Archives of General Psychiatry* 65(12):1429–1437, 2008. PMID: 19047530
- ²² Wechsler, H.; Dowdall, G.W.; Maenner, G.; et al. Changes in binge drinking and related problems among American college students between 1993 and 1997: Results of the Harvard School of Public Health College Alcohol Study. *Journal of American College Health* 47(2):57–68, 1998. PMID: 9782661
- ²³ Stratton, K.; Howe, C.; Battaglia, F.; Eds. *Fetal Alcohol Syndrome: Diagnosis, Epidemiology, Prevention, and Treatment*. Washington, DC: Institute of Medicine, National Academy Press, 1996.
- ²⁴ May, P.A.; Gossage, J.P.; Kalberg, W.O.; et al. Prevalence and epidemiologic characteristics of FASD from various research methods with an emphasis on recent in-school studies. *Developmental Disabilities Research Reviews* 15(3):176–192, 2009. PMID: 19731384
- ²⁵ May, P.A.; Keaster, C.; Bozeman, R.; et al. Prevalence and characteristics of fetal alcohol syndrome and partial fetal alcohol syndrome in a Rocky Mountain Region City. *Drug and Alcohol Dependence* 155:118–127, 2015. PMID: 26321671
- ²⁶ Estimated liver disease deaths include deaths with the underlying cause of death coded as alcoholic liver disease (K70), liver cirrhosis, unspecified (K74.3–K74.6, K76.0, K76.9), liver cancer (C22), or other liver diseases (K71, K72, K73, K74.0–K74.2, K75, and K76.1–K76.8). Number of deaths from Multiple Cause of Death Public-Use Data File, 2015 (<http://wonder.cdc.gov/mcd.html>). Alcohol-attributable fractions (AAFs) from CDC Alcohol-Related Disease Impact (http://nccd.cdc.gov/DPH_ARDI/Default/Default.aspx, Accessed 5/23/17). Prevalence of alcohol consumption from the National Survey on Drug Use and Health, 2015, for estimating indirect AAFs for liver cancer.
- ²⁷ Yoon, Y.H., and Chen, C.M. *Surveillance Report #105: Liver Cirrhosis Mortality in the United States: National, State, and Regional Trends, 2000–2013*. Bethesda, MD: National Institute on Alcohol Abuse and Alcoholism (NIAAA), 2016. Available at: <http://pubs.niaaa.nih.gov/publications/Surveillance105/Cirr13.pdf>. Accessed 11/7/16.
- ²⁸ Singal, A.K.; Guturu, P.; Hmoud, B.; et al. Evolving frequency and outcomes of liver transplantation based on etiology of liver disease. *Transplantation* 95(5):755–760, 2013. PMID: 23370710 (Please note: The “almost 1 in 3” figure aggregates the total number of transplants necessitated by alcoholic cirrhosis, alcoholic liver disease plus hepatitis C virus infection, and 40 percent of transplants necessitated by hepatocellular carcinoma.)
- ²⁹ National Cancer Institute. *Alcohol Consumption*, November 2015 update. Available at: <http://www.progressreport.cancer.gov/prevention/alcohol>. Accessed 9/19/16.
- ³⁰ National Institute on Alcohol Abuse and Alcoholism (NIAAA). NIAAA Council Approves Definition of Binge Drinking. *NIAAA Newsletter*, No. 3, Winter 2004. Available at: https://pubs.niaaa.nih.gov/publications/Newsletter/winter2004/Newsletter_Number3.pdf Accessed 9/19/16.
- ³¹ Substance Abuse and Mental Health Services Administration (SAMHSA). *Binge Drinking: Terminology and Patterns of Use*, 2016. Available at: <https://www.samhsa.gov/data/report/2017-methodological-summary-and-definitions>. Accessed 1/18/17.



National Institute
on Alcohol Abuse
and Alcoholism

NIH . . . Turning Discovery Into Health®

National Institute on Alcohol Abuse and Alcoholism

<https://www.niaaa.nih.gov> • 301-443-3860

Updated December 2019